


CyTRAK Orange™ in Live End-Point Imaging

Orange Fluorescent Live-Cell Permeant DNA and Cytoplasmic Dye

1. LIVE CELL NUCLEAR COUNTERSTAINING

BACKGROUND

In fluorescence-based microscopy it is normal to preserve cells or tissue sections with formaldehyde or similar fixative. This allows review of the samples at one's convenience and stops the loss or change of the desired biological readout. The fixative works by crosslinking with proteins and causes a mild permeabilisation of the lipid bilayer membranes. However, there are circumstances where fixation is detrimental, for example, if the target for analysis leaks into other sub-cellular compartments or leaks out of the cell completely on washing. Alternatively, the fixation process itself may produce artefacts. Similarly, fixation is inappropriate in assays to study drug-receptor interactions as cells are imaged after test antagonists and controls and then again after the reference agonist(s). The same would be true in assays to follow the internalization or processing of a surface-bound molecule (e.g. GPCRs). Nonetheless, such live-cell end-point assays, like those on formaldehyde-fixed samples, usually require a nuclear counterstain to demark the individual cells and, in the case of tissue, to give additional information on tissue morphology. Of course, on these samples, it is essential that the nuclear counterstain is fully cell permeant.



WHAT IS THE PROBLEM?

Cell impermeant DNA dyes are unusable as nuclear counterstains in such assays. The UV-excited dye Hoechst 33342 labels nuclei and is live cell permeant, however on some imaging platforms the coincidental detection of emission from Hoechst and GFP or fluorescein-derived functional probes mean these have to be illuminated sequentially slowing data acquisition and stretching time differences between reading the first and last samples which complicates the performance of live-end end-point assays. Additionally, the binding of Hoechst 33342 to dsDNA is time-sensitive and subject to clearance by MDR-phenotypes and ABCG2 pumps. Hoechst 33342 is supplied as hard compound and has to be solubilized initially with DMSO and cannot be stored in aqueous form, ready-to-use.

HOW DOES CyTRAK Orange™ HELP?

The orange fluorescing live cell permeant DNA probe CyTRAK Orange™ provides clear nuclear counterstaining in live cells. The bright signal also shows nuclear condensation and DNA fragmentation which can indicate deteriorating cell health. CyTRAK Orange™ gives a useful secondary cytoplasmic signal for further cellular segmentation. CyTRAK Orange™ is spectrally separated from commonly used fluorescent protein tags (e.g. CFP, GFP), UV- and red-excited and fluorescein-derived chromophores which eases assay design and allow simultaneous and rapid image acquisition. DNA binding is achieved in minutes, is temporally stable and unaffected by MDR phenotypes.

CyTRAK Orange™ is provided in an aqueous, ready-to-use solution. CyTRAK Orange™ is photo- and chemically-stable, widely compatible with physiological buffers. CyTRAK Orange™ is documented in live end-point analyses.

CyTRAK Orange™ Product Features:

- ❖ Orange-fluorescing cell-permeant dsDNA probe (blue/green excitation)
- ❖ rapidly and clearly labels all nuclei in live cells
- ❖ single-channel dual compartment (nucl:cyto) segmentation
- ❖ compatible with CFP, GFP, fluorescein-derived & red-excited chromophores
- ❖ water-soluble; ready-to-use from the fridge



Document Ref: CO5.APPNOTE LIVE ENDPOINT
Revision #: 002
Revision date: 09/06/26

BioStatus Limited
 56A Charnwood Road, Shepshed, Leicestershire LE12 9NP, United Kingdom
 T +44 1509 558 163 | E enquiry@biostatus.com | W www.biostatus.com
 For Research Use Only. © Copyright BioStatus. All Rights Reserved.