

CyGEL™ in Image-Based Screening


Thermoreversible Hydrogel Mountant for Non-adherent/Motile Cells

1. IMMOBILISING NON-ADHERENT CELLS/MICRO-TISSUES IN MICROPLATES

BACKGROUND

In cell-/tissue-based assays for high content screening (HCS, target-based or phenotypic) fluorescence-based imaging is the preferred approach when it is important to understand changes that occur in the context of morphology or of intracellular texture. This allows a number of parameters to be studied simultaneously: a disease-related readout such as a protein translocation, changes in cell numbers and morphology (sentinels for toxicity), a functional one such as mitochondrial health, cell:cell interactions or even organotypic relevance in the case of micro-tissues or model organisms.

In principle these assays can be performed on live adherent, non-adherent / suspension cells, micro-tissues and model organisms such as zebrafish.



WHAT IS THE PROBLEM?

The problem arises when non-adherent or motile objects need to be used in microplate formats for imaging since they are essentially mobile within the well. Non-adherent objects in a conventional culture medium or buffer often do not distribute evenly within the micro-well due to settling, rolling and clumping artifacts made worse by the 'panning' effects of movements of the microplate. Frequently non-adherent cells and motile organisms will concentrate in the centre or at the perimeter of the well. To overcome these issues suspension cells can be centrifuged onto the well surface and fixed or 'tacked' to the surface using adhesion reagents - all potentially introducing bias in the cells that are immobilized. Centrifugation may not be compatible with fragile micro-tissues and may not properly orientate model organisms.

HOW DOES CyGEL™ HELP?

An alternative provided by CyGEL™ is to immobilize these objects in an optically compatible hydrogel. CyGEL™ allows you to observe live objects for multiple illuminations, z-sections for 3D rendering or to re-visit a location without worrying about their rapid motility or movement in wells or on slides. It is not intended as a long-term growth matrix. CyGEL™ works as a thermoreversible hydrogel - *liquid when cold and a gel when warmed*. It is compatible with LIVE cells suspensions, micro-tissues and model organisms. The gel is formulated to be liquid when cool so can be conveniently stored at 4 °C and mixed with a cell preparation, or ready in wells for delivery of objects or used as an overlay in slides or wells. Upon warming beyond 22 °C to 37 °C, a CyGEL™ preparation rapidly transitions to a gel. This allows the operator to generate simple protocols for the capture and analysis of non-adherent or motile objects. CyGEL™ (PBS-formulated) is compatible with culture media, small molecules such as viability dyes, vital dyes and anaesthetics, fluorescence and standard microscopy optics. CyGEL™ does not disturb the imaging of fluorescent proteins. If required, CyGEL™ uniquely allows non-destructive recovery of live objects embedded in it.

CyGEL™ Product Features:

- ❖ convenient immobilisation of non-adherent cells and beads
- ❖ optically clear with low autofluorescence for visible range excitation
- ❖ controllable performance and rapid reversible transition from liquid to gel
- ❖ compatible with GFP and "in gel" fluorescent probes including DRAQ5™



Document Ref: CY1.APPNOTE IMAGE SCREEN
Revision #: 002
Revision date: 09/06/26

BioStatus Limited
 56A Charnwood Road, Shepshed, Leicestershire LE12 9NP, United Kingdom
 T +44 1509 558 163 | E enquiry@biostatus.com | W www.biostatus.com
 For Research Use Only. © Copyright BioStatus. All Rights Reserved.